

ANTIMICROBIAL ACTIVITY OF FRACTIONS AND SUBFRACTIONS OF *ELAEAGIA UTILIS* AGAINST MICROORGANISMS OF IMPORTANCE IN DENTAL CARIES

Jennyfer Aldana^{1,2}, Nohemí Téllez², Fredy Gamboa^{1,3}

¹Dental Research Centre Group, School of Dentistry, Pontificia Universidad Javeriana. Bogotá, Colombia.

²Department of Chemistry, Phytochemistry Group, Pontificia Universidad Javeriana. Bogotá, Colombia.

³Department of Microbiology, School of Sciences, Pontificia Universidad Javeriana. Bogotá, Colombia.

ABSTRACT

Dental caries is a multifactorial infectious disease that leads to the destruction of dental hard tissue. The main goal of research into medicinal plants is to seek compounds with antimicrobial activity for subsequent use in prevention strategies and control of infectious diseases. The aim of this study was to evaluate the antimicrobial activity of fractions and subfractions obtained from *Elaeagia utilis* against *Streptococcus mutans*, *Streptococcus sobrinus* and *Lactobacillus acidophilus*. The plant material was collected in the town of Alban (Cundinamarca, Colombia), which is located at an altitude of 2245 meters above sea level. Two extracts were obtained by cold maceration of *E. utilis* leaves in (a) petroleum ether extract and (b) ethanol extract. Fractions were obtained from the petroleum ether extract by column vacuum chromatography, and from the ethanol extract by continuous liquid / liquid partitioning. The antimicrobial activity of frac-

tions and subfractions was evaluated by the well diffusion method. At a concentration of 10 mg/well, several fractions from both extracts showed antimicrobial activity against *S. mutans*, *S. sobrinus* and *L. acidophilus*. Among the ethanol extract fractions, the dichloromethane fraction had notably greater antimicrobial activity. It was sub-partitioned, yielding three subfractions with inhibitory activity, of which the most active was MeOH: H₂O (Bp) with minimum inhibitory concentration 0.1 mg/well on the 3 study bacteria. Terpenes, sesquiterpenlactones and simple phenolic compounds were identified in it. In conclusion, this study shows the antimicrobial potential of fractions and subfractions obtained from extracts of *E. utilis* leaves against bacteria that are important in dental caries.

Keywords: Plants, medicinal, Dental caries, *Streptococcus mutans*, *Streptococcus sobrinus*, *Lactobacillus acidophilus*.

ACTIVIDAD ANTIMICROBIANA DE FRACCIONES Y SUBFRACCIONES DE *ELAEAGIA UTILIS* SOBRE MICROORGANISMOS DE IMPORTANCIA EN CARIES DENTAL

RESUMEN

La caries dental es una enfermedad infecciosa multifactorial que conduce a la destrucción del tejido duro dental. El principal objetivo de la investigación en plantas medicinales es la búsqueda de compuestos con actividad antimicrobiana para su posterior uso en estrategias de prevención o control de enfermedades infecciosas. El objetivo de este estudio fue evaluar la actividad antimicrobiana de fracciones y subfracciones obtenidas de la planta *Elaeagia utilis* contra *Streptococcus mutans*, *Streptococcus sobrinus* y *Lactobacillus acidophilus*. El material vegetal fue colectado en la ciudad de Albán (Cundinamarca-Colombia) situada a una altitud de 2245 metros sobre el nivel del mar. Mediante el método de maceración en frío de hojas de *E. utilis* se obtuvieron dos extractos, uno en éter de petróleo y otro en etanol. Del extracto etéreo se obtuvieron fracciones mediante cromatografía en columna al vacío y al extracto etanólico se le realizó fraccionamiento líquido/líquido continuo. La evaluación de la actividad antimicrobiana de las fracciones y

subfracciones se realizó por el método de difusión en pozo. A una concentración de 10 mg/pozo, múltiples fracciones obtenidas de los dos extractos presentaron actividad antimicrobiana sobre *S. mutans*, *S. sobrinus* y *L. acidophilus*. De las fracciones del extracto etanólico se destaca la fracción diclorometano, por presentar mayor actividad antimicrobiana, razón por lo cual se subfracciona y se obtienen tres subfracciones con actividad inhibitoria. La subfracción más activa fue MeOH:H₂O (Bp) con una concentración mínima inhibitoria de 0.1 mg/pozo sobre las 3 bacterias en estudio. En esta subfracción se determinaron terpenos, sesquiterpenlactonas y compuestos fenólicos simples. En conclusión, en este estudio se presenta el potencial antimicrobiano de fracciones y subfracciones obtenidas de extractos de hojas de *E. utilis* contra microorganismos de importancia en caries dental.

Palabras claves: Plantas medicinales, caries dental, *S. mutans*, *S. sobrinus*, *L. acidophilus*.

INTRODUCTION

Dental caries is one of the most common chronic, multifactorial, transmittable infectious diseases in the world¹. *Streptococcus mutans* and to a lesser extent, *Streptococcus sobrinus*, *Streptococcus gor-*

donii, and species of *Lactobacillus* and *Actinomyces* are the primary microorganisms related to the development and progression of dental caries¹⁻³. Recognizing their importance in the initiation and progression of caries leads to designing meas-

ures aimed at eliminating or reducing them in the oral cavity⁴.

Plant species have been widely used around the world as a source of traditional medicines for treating diseases.⁵ The primary aim of research into medicinal plants is to identify plants with pharmacological activity in order to discover new substances with antimicrobial activity, which through various chemical procedures can be made into medications to control or prevent infectious diseases⁶. Many substances from various plant families have antimicrobial activity useful for oral health.⁶⁻¹⁰ Allicine, from *Allium sativum* (Amaryllidaceae)⁷, macelignan from *Myristica fragrans* (Myristicaceae)⁸, bakuchiol from *Psoralea coryfolia* (Leguminosae)⁹, and isopanduratin A from *Kaempferia pandurata* (Zingiberaceae)¹⁰, are examples of molecules of natural origin with antimicrobial activity against *S. mutans*, *S. sobrinus*, *S. salivarius* and other microorganisms that are important in the oral cavity.

The family Rubiaceae, one of the 5 plant families with greatest ecological and taxonomical diversity in the world, has a large number of genera and species with special distribution in tropical Andean rainforests¹¹. In Colombia, it is represented by 105 native genera including over 960 species, which grow mainly in the Andean, Amazonian and Chocó biogeographic regions¹¹.

Rubiaceae family members which are known to have antimicrobial potential against microorganisms that are important in oral infections are *Uncaria tomentosa* against *S. mutans*¹² and fractions obtained from *Isertia laevis* leaves against *S. mutans* and *S. sobrinus*¹³. This draws attention to the need for further research to foster the discovery of molecules or substances that could be extracted from plants in this family and may help eliminate or reduce cariogenic microorganisms.

Species of the genus *Elaeagia* (family Rubiaceae) are distributed throughout the Colombian Andes region, at elevations of 100 to 2,600 meters above sea level¹¹, and are known as “Barniz” and “Mopamopa”^{11,14}. There is currently no information on evaluation of the antimicrobial activity of *Elaeagia utilis* against microorganisms that are important to dental caries and oral health. The aim of this study was to assess the antimicrobial activity of fractions and subfractions obtained from *Elaeagia utilis* leaves against *S. mutans*, *S. sobrinus* and *L. acidophilus*.

MATERIALS AND METHODS

1. Obtaining and processing plant material

The plant material was collected in the municipality of Albán (Cundinamarca, Colombia), at Padre Luna's Foundation “Granjas Infantiles” (Children's Farms). The specimen was taxonomically determined at the Herbarium of Javeriana University as *Elaeagia utilis* (Goudot) Wedd, of the family Rubiaceae, collection number HPUJ 27413. The leaves were dried at room temperature, then ground and powdered using a chopper. Cold maceration of the dried plant material (960 g) in petroleum ether (petrol) for 72 hours yielded an extract in petrol. The plant material filtered out of the first extraction was placed in cold maceration in ethanol (EtOH), and yielded an ethanol extract.

2. Obtaining fractions

A. Fractions from the petroleum ether extract

One gram of the extract in petrol was passed through a chromatography column under vacuum with a stationary phase composed of silica 60-H (0.063-0.200mm; Merck, Germany), in a proportion of 30:1 for stationary phase:sample. It was eluted with solvents of different polarities: Petrol, Petrol: Dichloromethane (CH₂Cl₂) (1:1), CH₂Cl₂, CH₂Cl₂:ethyl acetate (AcOEt) (1:1), AcOEt, AcOEt:EtOH (1:1) and EtOH.

B. Fractions from the ethanol extract

The ethanol extract was subject to continuous liquid/liquid partitioning (CLLP) and yielded four fractions: Petrol, CH₂Cl₂, AcOEt and Butanol (BuOH). Each fraction was concentrated at 40°C, under reduced pressure in a rotary evaporator (Buchi B-169, Vacuum-System; Germany) until it was dry. The fraction with greatest antimicrobial activity was sub-partitioned by column vacuum chromatography with stationary phase RP-18 (40-63 μm; Merck, Germany) and eluted with methanol:water (MeOH:H₂O), MeOH and MeOH:CH₂Cl₂.

3. Methods for chemical characterization of active subfractions

Qualitative chemical tests were performed on the subfraction with greatest antimicrobial activity to determine types of secondary metabolites. Then it was partitioned by gas chromatography mass spectrometry (GC-MS) in a chromatograph (Agilent Technology 6850 series II) connected to an electron

impact mass spectrometer (70eV) model Agilent MS 5975B. The device has a fused-silica capillary column with 5% polydimethylsiloxane stationary phase (30 m long, 0.25 mm diameter and 0.25 μm phase thickness), using as carrier gas 99.995%, Aga Fano, S.A, grade 5, with a constant flow of 1 mL/min. The sample was dissolved in methanol ($1\text{mg}\cdot\text{mL}^{-1}$) and $1\mu\text{L}$ was injected in Split 15:1 mode. The initial temperature was 80°C , for 2 minutes, increasing by $10^{\circ}\text{C}/\text{min}$ up to 280°C sustained for 5 minutes. Data from the 7th edition of the Wiley Mass Spectra library were compared to those obtained from the sample for identification. Spectra with matches better than 90% were considered to provide adequate identification.

4. Evaluation of the antimicrobial activity of fractions and subfractions

A. Study strains

Antimicrobial activity was evaluated on three reference strains: *S. mutans* ATCC 25175, *S. sobrinus* CIO 428 and *L. acidophilus* ATCC 4365, which had been preserved by freezing at -70°C at the Dental Research Center at Javeriana University. In order to reconstitute them and confirm viability, $20\mu\text{L}$ from the preservation vials were thawed and cultured in brain heart infusion (BHI) broth for 4 hours at 37°C in an anaerobic atmosphere ($\text{H}_2:\text{CO}_2:\text{N}_2$; 10:10:80). Bacteria grown in the BHI broth were plated on BHI agar and incubated for 16 hours at 37°C in an anaerobic atmosphere ($\text{H}_2:\text{CO}_2:\text{N}_2$; 10:10:80). Pure, viable colonies of *S. mutans* ATCC 25175, *S. sobrinus* CIO 428 and *L. acidophilus* ATCC 4365 were reconfirmed using Gram stain and biochemical tests.

B. Well diffusion method

The antimicrobial activity of the *E. utilis* extracts, fractions and subfractions against bacteria was evaluated using the well diffusion technique described by Dobner et al.¹⁵ A suspension was prepared from each culture of pure bacteria and adjusted by turbidimetry to a 0.5 McFarland standard. Then $100\mu\text{L}$ of the suspension were added to 20 mL liquid Mueller Hinton agar, mixed and poured into Petri dishes. Twenty minutes after it had solidified, a sterilized glass Pasteur pipette was used to make wells 0.5 cm in diameter on the agar. Fifty μL of the extract, fraction or subfraction dissolved in dimethyl sulfoxide (DMSO),

were placed individually into each well. Fifty μL of Vancomycin at a concentration of $150\mu\text{g}/\text{mL}$ were used as a positive control and $50\mu\text{L}$ DMSO as a negative control. The dishes were immediately incubated at 37°C for 24 hours. Tetrazolium salt (MTT; $2.5\text{mg}/\text{mL}$ aqueous solution) was added to the surface to reveal bacterial viability¹⁶, and the dishes were left to incubate for another six hours. After incubation, the zones of inhibition produced by the fractions and/or subfractions were measured and the minimum inhibitory concentration (MIC; the lowest concentration producing a zone of inhibition of at least 6 mm) was determined. Each test was performed in triplicate and the average reported in mm.

4. Evaluation of the antimicrobial effect of the active subfraction by bioautography

The subfraction which was shown to have antimicrobial activity according to the well diffusion method was also evaluated using the bioautographic method described by Cos et al.¹⁷ and Valgas et al.¹⁸ It was applied on chromatographic plates 7.5 cm long and 2.5 cm wide with silica gel stationary phase (60 μm F254, Merck, Germany). The mobile phase was $\text{CH}_2\text{Cl}_2:\text{MeOH}$ (9:1). The chromatographic plates were sterilized by exposure to UV radiation (wavelength 260nm) for 30 minutes. Suspensions of each of the 3 study bacteria were prepared and adjusted to a 0.5 McFarland standard. One hundred μL of each suspension was added to 10 mL liquid Mueller Hinton agar, mixed and poured onto the chromatography plate in the Petri dish. The chromatography plates were incubated for 24 hours at 37°C , after which MTT was added and they were returned to incubation at 37°C for another 6 hours. Rf (distance travelled by solute / distance travelled by solvent) was measured in the zones of inhibition (colourless areas).

RESULTS

Antimicrobial activity of the fractions from the extract in petroleum ether

Table 1 shows the antimicrobial activity of the extract in petrol and the fractions derived from it against the three study bacteria with 10 mg/well. Only two fractions (Petrol: CH_2Cl_2 and CH_2Cl_2) had no antimicrobial activity against the bacteria. The other 4 fractions produced zones of inhibition ranging from 7 to 15 mm.

Table 1: Antimicrobial activity of the Petroleum Ether extract and its subfractions against *S. mutans* ATCC 25175, *S. sobrinus* CIO 428 and *L. acidophilus* ATCC 4365 at a concentration of 10mg/well. Inhibitory activity (zones of inhibition) in mm.

Total extract and fractions		<i>S. mutans</i> ATCC 25175	<i>S. sobrinus</i> CIO 428	<i>L. acidophilus</i> ATCC 4365
Total extract	Petroleum Ether extract	9.5	9	9
Fractions	Petrol:CH ₂ Cl ₂	0	0	0
	CH ₂ Cl ₂	0	0	0
	CH ₂ Cl ₂ :AcOEt	11	9	9
	AcOEt	8.5	9	7
	AcOEt:EtOH	15	8.5	8
	EtOH	11	8.5	8.5
Positive control	Vancomycin (150 µg/mL)	19	17	19
Negative control	DMSO	0	0	0

Antimicrobial activity of the fractions obtained by CLLP from the extract in ethanol


Antimicrobial activity was tested for the Petrol, CH₂Cl₂, AcOEt and BuOH fractions obtained by liquid/liquid partitioning of the ethanol extract, plus some precipitates which appeared upon partitioning with AcOEt and BuOH, which were named AcOEt(p) and BuOH(p). Fig. 1 shows the antimicrobial activity of the fractions and precipitates on the 3 study bacteria. The dichloromethane fraction stands out with a larger zone of inhibition.

Antimicrobial activity of the subfractions yielded by the active CH₂Cl₂ fraction from the extract in ethanol

Because the CH₂Cl₂ fraction in the ethanol extract had outstanding antimicrobial activity, it was separated using CVC with stationary phase RP-18, 30:1 (Station-

ary Phase:Sample), and mobile phase MeOH:H₂O (10:1), MeOH and MeOH:CH₂Cl₂ (9:1). This procedure yielded seven subfractions: MeOH:H₂O (A)(0.5 mg), MeOH:H₂O (B)(1.52 mg), MeOH:H₂O (C)(0.2 mg), MeOH:H₂O (D)(0.08 mg), MeOH:H₂O (E)(0.05 mg), MeOH:H₂O (F)(0.15 mg) and MeOH:CH₂Cl₂ (0.63 mg). Subfraction MeOH:H₂O (B) had a precipitate, which was separated from the supernatant and named MeOH:H₂O (Bp). Fig. 2 shows the antimicrobial activity of the eight subfractions obtained from the dichloromethane fraction. Subfractions MeOH:H₂O A, MeOH:H₂O B and MeOH:H₂O Bp have outstanding inhibitory activity. Subfraction MeOH:H₂O (Bp) at 10 mg/well had the greatest biological activity against the bacterial strains, so we decided to evaluate it at lower concentrations, and found it was active to a concentration of 0.1 mg/well against the three study bacteria (Fig. 3).

Fig. 1: A: Antimicrobial action of the fractions and precipitates (p) at 10 mg/well obtained by continuous liquid/liquid partitioning (CLLP) of the ethanol extract. B: Zone of inhibition (17 mm) of fraction CH₂Cl₂ against *S. sobrinus* CIO 428. Vancomycin and dimethyl sulfoxide were used, respectively, as positive control -VAN (C+) and negative control-DMSO (C-).






Fig. 2: A: Antimicrobial activity of the subfractions (10 mg/well) obtained from the CH₂Cl₂ fraction of the ethanol extract. B: Thin layer chromatography (stationary phase: RP-18, mobile phase: MeOH:H₂O; 10:1); of subfractions MeOH:H₂O B (1) and MeOH:H₂O B (2).




Fig. 3: A: Antimicrobial activity of subfraction MeOH:H₂O (Bp) at three different concentrations. B: Zones of inhibition on *S. mutans* ATCC 25175 produced by subfraction MeOH:H₂O (Bp) at concentrations of 1mg/well, 0.5mg/well and 0.1mg/well.

Antimicrobial activity of the active subfraction by bioautography

Fig. 4 shows the antimicrobial activity of subfraction MeOH:H₂O (Bp) against the three study bacteria using the bioautographic technique. The antimicrobial activity of the compounds was located at an Rf zone located between 0.44 and 0.49.

Chemical study of the active subfraction

The following qualitative chemical tests were performed: Baljet, Lieberman-Burchard, Salkowski,

ammonium molybdate, iron (III) chloride, Dragendorff, foam test, anthrone, ferric hydroxamate. The Baljet, Salkowski, ammonium molybdate, foam and ferric hydroxamate tests were positive, indicating presence of diterpenes, steroids and saponins. The others were negative.

Table 2 shows the compounds identified by GC-MS in the active subfraction MeOH:H₂O (Bp). The most outstanding compounds found in the mixture were simple phenolic compounds, benzene derivatives and hydrocarbons.

Fig. 4: Antimicrobial activity of subfraction MeOH:H₂O (Bp) by bioautography (Chromatographic plate: stationary phase silica gel and mobile phase CH₂Cl₂:MeOH, 9:1, with MTT reagent) on *S. mutans* ATCC 25175 (A) and *L. acidophilus* ATCC 4365 (B). C shows the compounds responsible for the inhibition under UV light (wavelength 254 nm) with Rf values.




Table 2: Compounds identified in subfraction MeOH:H₂O (Bp) by GC-MS (Agilent Technology 6850 series II, connected to an electron impact mass spectrometer (70eV) Agilent MS 5975B).

Retention time (seconds)	Name of compound	Area covered (%)	Nature of the Compound
5.885	Benzyl alcohol	1.793	Benzene derivative
7.475	Benzoic acid	1.409	Benzene derivative
8.491	4-vinylphenol	1.186	Simple phenolic compound
8.504	2,3-dihydrobenzofuran	1.186	Aromatic hydrocarbon
9.807	2-methoxy-4-vinylphenol	2.427	Simple phenolic compound
11.775	3-methoxy-4-hydroxybenzaldehyde (vanillin)	0.430	Simple phenolic compound
13.483	2,5-dimethyl-(3-methoxymethyl)-p-benzoquinone	1.116	Benzene derivative
13.732	2,6-dimethoxy-4-(2-propenyl) phenol	0.553	Simple phenolic compound
16.483	1,Z-5,E-7-dodecatrien	2.344	Hydrocarbon
16.662	Loliolide	0.808	Sesquiterpene lactone
16.973	Benzoic acid, 2-hydroxy-, phenyl methyl ester	1.846	Benzene derivative
16.014	4-((1e)-3-hydroxy-1-propenyl)-2- methoxyphenol	9.443	Simple phenolic compound

DISCUSSION

Colombian flora is widely known, and is considered to be a potential source of products with pharmacological activity.¹⁹ Many substances obtained from plant species have been evaluated against pathogenic microorganisms, and their antimicrobial activity has been proven and/or reconfirmed.^{7-10,17,19} Natural compounds with antimicrobial activity are the basis for a line of research that seeks to discover structural and functional components, called active principles, most of which are secondary

metabolites.^{6,12,13,19-21} Research into these compounds is a strategic route for the development of efficacious affordable drugs which can be used for treating diseases that are important to the public.⁶⁻¹⁰ Prior research into plant species of the Rubiaceae family reports antimicrobial activity against different microorganisms.^{19,21-23} The ethanol extract from *Cinchona officinalis* had antimicrobial activity against *S. aureus*, *Bacillus cereus* and β -hemolytic *Streptococcus*.¹⁹ Similarly, the ethanol extract from *Uncaria tomentosa* bark inhibited the growth of 6

bacteria, of which the most outstanding are *Bacillus subtilis*, *Enterococcus faecalis*, *Escherichia coli*, *Staphylococcus aureus*.²¹ Another study on *Uncaria tomentosa* aqueous extracts showed antimicrobial activity against *S. aureus* and *Candida albicans*.²² Studies using *in vitro* models report that *Coffea arabica* aqueous extracts reduce the adherence of *S. mutans* to dental enamel and dentin.²³

The microbiological evaluation of *E. utilis* leaves showed that both the petroleum ether extract and 4 of the 6 fractions derived from it had antimicrobial activity against the 3 study bacteria. Similarly, 5 of the 6 fractions obtained from the ethanol extract by liquid/liquid extraction had antimicrobial activity against *S. mutans*, *S. sobrinus* and *L. acidophilus* at a concentration of 10 mg/well. Of these, the dichloromethane fraction was outstanding because it produced larger zones of inhibition against all 3 microorganisms, which is why it was sub-partitioned.

Three of the 8 subfractions obtained from the dichloromethane fraction had inhibitory action against the 3 study bacteria. Of these three subfractions, MeOH:H₂O (Bp) was outstanding because it produced the largest zones of inhibition (15-20 mm). The fact that the activity of subfraction MeOH:H₂O (Bp) was greater than that of subfraction MeOH:H₂O (B) may be due to the fact that the former has a precipitation process leading to the separation of additional secondary metabolites which may act as interference with biological activity. Because subfraction MeOH:H₂O (Bp) had less mixture of components and higher biological activity, it was selected for microbiological assays at lower concentrations and determination of MCI, which was found to be 0.1 mg/well.

In the bioautography, subfraction MeOH:H₂O (Bp) showed antimicrobial activity against all 3 microorganisms and when vanillin was used as a reagent, it showed as a uniform yellow patch; nevertheless, exposure to UV light revealed two patches located very close together, with Rf 0.44 and 0.49, suggesting that the compounds in it are highly complex.

The qualitative chemical study performed on subfraction MeOH:H₂O (Bp), primarily determined presence

of terpenes, sesquiterpene lactones and saponins. These findings agree with De Rosa et al.²⁴ and Zhao et al.²⁵, who report the presence of triterpenic saponins in other Rubiaceae species. Moreover, Kloucek et al.²¹ report that triterpenes, are major components in mixtures with antimicrobial activity obtained from species from the family Rubiaceae.

GC-MS analysis of subfraction MeOH:H₂O (Bp) showed presence of various classes of compounds, among which simple phenols and benzene derivatives are outstanding. Cowan²⁶ reports that simple phenols, phenolic acids and quinones are the main components of plant origin that have antimicrobial activity. Gopalakrishnan et al.²⁷ report antimicrobial activity of 4-((1e)-3-hydroxy-1-propenyl)-2-methoxyphenol, a compound which is present in the subfraction of this study. Moreover, Friedman et al.²⁸ report that the commercial compounds 3-methoxy-4-hydroxybenzaldehyde and benzoic acid, 2-hydroxy-phenyl methyl ester, which are also present in subfraction MeOH:H₂O (Bp), have antimicrobial activity against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Escherichia coli*. With regard to the compounds 4-vinylphenol and 2-methoxy-4-vinylphenol found in the mixture of subfraction MeOH:H₂O (Bp), other studies report that they are major components in leaf extracts from the species,²⁹ while in acetone extract from *Rumex vesicarius* (Polygonaceae) leaves with activity against *E. coli* and *C. albicans*, one of the main components is 2-methoxy-4-vinylphenol.³⁰

The results of this study show the potential of fractions and subfractions obtained from *E. utilis* leaf extracts as sources of various compounds with antimicrobial activity against bacteria that are important in dental caries. In this regard, further studies are needed to isolate, characterize and identify substances present in active fractions and subfractions, and to determine their antibacterial activity against a wide range of microorganisms that are important in other oral infections. In the future, these substances may be used in toothpastes, mouth rinses and other products for oral hygiene and health.

ACKNOWLEDGMENTS

The authors would like to thank the Academic Vice-director's office of Javeriana University for financing the research project (number 002286).

CORRESPONDENCE

Dr. Fredy Gamboa, Ph.D.
Departamento de Microbiología (Facultad de Ciencias)
Pontificia Universidad Javeriana
Carrera 7 No. 40-62- Bogotá- Colombia
E-mail: gamboa@javeriana.edu.co

REFERENCES

1. Loesche WJ. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev* 1986;50:353-380.
2. Belda-Ferre P, Alcaraz LD, Cabrera-Rubio R, Romero H, Simón-Soro A, Pignatelli M, Mira A. The oral metagenome in health and disease. *ISME J* 2012;6:46-56.
3. Aas JA, Griffen AL, Dardis SR, Lee AM, Olsen I, Dewhirst FE, Leys EJ, Paster BJ. Bacteria of dental caries in primary and permanent teeth in children and young adults. *J Clin Microbiol* 2008;46:1407-1417.
4. Azevedo MS, van de Sande FH, Romano AR, Cenci MS. Microcosm biofilms originating from children with different caries experience have similar cariogenicity under successive sucrose challenges. *Caries Res* 2011;45:510-517.
5. Hamburger M, Hostettmann K. Bioactivity in plants: The link between phytochemistry and medicine. *Phytochemistry* 1991;30:3864-3874.
6. Sharapin N: Fundamentos de tecnología de productos fitoterapéuticos. Bogotá, Colombia: Ed. Roberto Pinzón, Área de ciencia y tecnología del Convenio Andrés Bello, 2000:225-251
7. Bakri IM, Douglas CW. Inhibitory effect of garlic extract on oral bacteria. *Arch Oral Biol* 2005;50:645-651.
8. Chung JY, Choo JH, Lee MH, Hwang JK. Anticariogenic activity of macelignan isolated from *Myristica fragrans* (nutmeg) against *Streptococcus mutans*. *Phytomedicine* 2006; 13:261-266.
9. Katsura H, Tsukiyama RI, Suzuki A, Kobayashi M. In vitro antimicrobial activities of bakuchiol against oral microorganisms. *Antimicrob Agents Chemother* 2001;45:3009-3013.
10. Hwang JK, Chung JY, Baek NI, Park JH. Isopanduratin A from *Kaempferia pandurata* as an active antibacterial agent against cariogenic *Streptococcus mutans*. *Int J Antimicrob Agents* 2004;23:377-381.
11. Mendoza H, Ramírez B, Jiménez L: Rubiaceae de Colombia Guía ilustrada de géneros. Bogotá, Colombia: Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, 2004:1-351.
12. Ccahuana-Vasquez RA, Santos SS, Koga-Ito CY, Jorge AO. Antimicrobial activity of *Uncaria tomentosa* against oral human pathogens. *Braz Oral Res* 2007;21:46-50.
13. Tellez N, Tellez M, Perdomo M, Alvarado A, Gamboa F. Anticariogenic activity of the active fraction from *Iserbia laevis* against *S. mutans* and *S. sobrinus*: comparison of two extraction methods. *Acta Odontol Latinoam* 2010; 23:188-195.
14. Jiménez L. Lista de las colecciones colombianas de Rubiaceae depositadas en el Herbario Nacional Colombiano (COL). *Caldasia* 2002;24:41-64.
15. Dobner MJ, Schwaiger S, Jenewein IH, Stuppner H. Antibacterial activity of *Leontopodium alpinum* (Edelweiss). *J Ethnopharmacol* 2003;89:301-303.
16. Thom SM, Horobin RW, Seidler E, Barer MR. Factors affecting the selection and use of tetrazolium salts as cytochemical indicators of microbial viability and activity. *J Appl Bacteriol* 1993;74:433-443.
17. Cos P, Vlietinck AJ, Berghe DV, Maes L. Anti-infective potential of natural products: how to develop a stronger in vitro 'proof-of-concept'. *J Ethnopharmacol* 2006;106:290-302.
18. Valgas C, Machado de Souza S, Smânia EFA, Smânia Jr A. Screening methods to determine antibacterial activity of natural products. *Braz J Microbiol* 2007;38:369-380.
19. Rojas JJ, Ochoa VJ, Ocampo SA, Muñoz JF. Screening for antimicrobial activity of ten medicinal plants used in Colombian folkloric medicine: a possible alternative in the treatment of non-nosocomial infections. *BMC Complement Altern Med* 2006;6:2.
20. Khan MR, Omoloso AD. Antimicrobial activity of extractives of *Sarcocephalus coadunatus*. *Fitoterapia* 2003;74:695-698.
21. Kloucek P, Polesny Z, Svobodova B, Vlkova E, Kokoska L. Antibacterial screening of some Peruvian medicinal plants used in Calleria District. *J. Ethnopharmacol* 2005; 99:309-312.
22. Herrera DR, Tay LY, Rezende EC, Kozłowski VA, Santos EB. In vitro antimicrobial activity of phytotherapeutic *Uncaria tomentosa* against endodontic pathogens. *J Oral Sci* 2010;52:473-476.
23. De Oliveira LD, da Silva BEH, Landucci LF, Koga-Ito CY, Olavo A, Jorge AOC. Effects of *Coffea arabica* on *Streptococcus mutans* adherence to dental enamel and dentine. *Braz J Oral Sci* 2007;6:1438-1441.
24. De Rosa S, Iodice C, Mitova M, Handjieva N, Popov S, Anchev M. Triterpene saponins and iridoid glucosides from *Galium rivale*. *Phytochemistry* 2000;54:751-756.
25. Zhao WM, Wolfender JL, Hostettman K, Cheng KF, Xu RS, Qin GW. Triterpenes and triterpenoid saponins from *Mussaenda pubescens*. *Phytochemistry* 1997;45:1073-1078.
26. Cowan MM. Plant products as antimicrobial agents. *Clin Microbiol Rev* 1999;12:564-582.
27. Gopalakrishnan S, Vadivel E. GC-MS analysis of some bioactive constituents of *Mussaenda frondosa* Linn. *Int J Pharm Bio Sci* 2011;2:313-320.
28. Friedman M, Henika PR, Mandrell RE. Antibacterial activities of phenolic benzaldehydes and benzoic acids against *Campylobacter jejuni*, *Escherichia coli*, *Listeria monocytogenes* and *Salmonella enterica*. *J Food Prot* 2003;66: 1811-1821.
29. Chuyen NV, Kurata T, Kato H, Fujimaki M. Antimicrobial activity of kumazasa (*Sasa albo-marginata*). *Agric Biol Chem* 1982;46:971-978.
30. Ara I, Bukhari NA, Solaiman D, Bakir MA. Antimicrobial effect of local medicinal plant extracts in the Kingdom of Saudi Arabia and search for their metabolites by gas chromatography-mass spectrometric (GC-MS) analysis. *J Med Plants Res* 2012;6:5688-5694.